WHAT IS CLAIMED IS:

 A method for neutralizing the anti-coagulant effect of heparin comprising administering to a subject an effective amount of a heparin binding BPI protein product.

5

- A method of inhibiting angiogenesis comprising administering to a subject an amount of a BPI protein product effective to inhibit angiogenesis.
- A method for neutralizing the anti-coagulant effect of heparin
 in a fluid sample comprising contacting said sample with an effective amount of heparin binding BPI protein product.
 - The method of claim 2 wherein the angiogenesis inhibited is associated with ocular retinopathy.

15

30

- A method for inhibiting endothelial cell proliferation comprising administering to a subject an amount of a BPI protein product effective to inhibit proliferation.
- A method of treating endometriosis comprising administering an amount of BPI to the endometrium effective to inhibit endothelial cell proliferation.
- A method for contraception comprising administering to a
 subject an amount of a BPI protein product to the uterine lining effective to prevent endothelial cell proliferation associated with implantation of a fertilized ovum.
 - A method for inhibiting malignant tumor cell proliferation comprising administering to a subject an amount of a BPI protein product effective to inhibit proliferation.

10

20

30

- The method of claim 8 wherein the malignant tumor is Kaposi's sarcoma.
- A method for treating a chronic inflammatory disease state
 comprising administering to a subject an amount of a BPI protein product effective to reduce inflammation.
 - 11. The method of claim 10 wherein the chronic inflammatory disease is arthritis.
 - The method of claim 11 wherein the arthritic inflammatory disease state is rheumatoid arthritis.
- The method of claim 11 wherein the arthritic inflammatory
 disease state treated is reactive arthritis.
 - 14. The method of claims 1, 2, 5, 6, 7, 8 and 10 wherein the BPI protein product is a 23-25 kD amino-terminal fragment of bactericidal/permeability-increasing protein.
 - 15. A method of inhibiting antiogenesis comprising administering to a subject an amount of a BPI protein product effective to inhibit angiogenesis.
- The method of claim 15 wherein the angiogenesis inhibited is
 associated with ocular retinopathy.
 - 17. A method for inhibiting endothelial cell proliferation comprising administering to a subject an amount of a bactericidal/permeability increasing (BPI) protein product effective to inhibit proliferation of endothelial cells.

10

15

- 18. A method of treating endometriosis comprising administering an amount of BPI to the endometrium effective to inhibit endothelial cell proliferation.
- 5 19. A method for contraception comprising administering to a subject an amount of a BPI protein product to the uterine lining effective to provent endothelial cell proliferation associated with implantation of a fertilized ovum.
 - 20. A method of inhibiting malignant tumor cell proliferation comprising administering to a subject an amount of a BPI protein product effective to inhibit proliferation.
 - The method of claim 20 wherein the malignant tumor is Kaposi's sarcoma.
 - 22. The method of claim 17 wherein the BPI protein product is an approximately 21 to 25 kD amino-terminal fragment of BPI protein.